

# The Resolved Stellar Population of the Post-Starburst Galaxy NGC 1569<sup>1</sup>

Laura Greggio<sup>2,3</sup>, Monica Tosi<sup>4</sup>, Mark Clampin<sup>5</sup>, Guido De Marchi<sup>6</sup>, Claus Leitherer<sup>5</sup>,  
Antonella Nota<sup>5,7</sup>, and Marco Sirianni<sup>8</sup>

<sup>2</sup> Università di Bologna, Dipartimento di Astronomia, Via Zamboni 33, I-40126 Bologna,  
Italy

e-mail: greggio@astbo3.bo.astro.it

<sup>3</sup> Universitaets Sternwarte Muenchen, Scheinerstrasse 1, D-81679 Muenchen, Germany

e-mail: greggio@usm.uni-muenchen.de

<sup>4</sup> Osservatorio Astronomico di Bologna, Via Zamboni 33, I-40126 Bologna, Italy

e-mail: tosi@astbo3.bo.astro.it

<sup>5</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

e-mail: clampin@stsci.edu, leitherer@stsci.edu, nota@stsci.edu

<sup>6</sup> European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching, Germany

e-mail: demarchi@eso.org

<sup>7</sup> Affiliated with the Astrophysics Division, Space Science Department of the European  
Space Agency

<sup>8</sup> Università di Padova, Dipartimento di Astronomia, Vicolo dell'Osservatorio 5, I-40100  
Padova, Italy

e-mail: sirianni@astrpd.pd.astro.it

---

<sup>1</sup>Based on observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by AURA for NASA under contract NAS5-26555

Received \_\_\_\_\_;    accepted \_\_\_\_\_

## ABSTRACT

We present WFPC2–HST photometry of the resolved stellar population in the post-starburst galaxy NGC 1569. The color-magnitude diagram (CMD) derived in the F439W and F555W photometric bands contains  $\sim 2800$  stars with photometric error  $\leq 0.2$  mag down to  $m_{F439}$ ,  $m_{F555} \simeq 26$ , and is complete for  $m_{F555} \lesssim 23$ . Adopting the literature distance modulus and reddening, our CMD samples stars more massive than  $\sim 4 M_{\odot}$ , allowing us to study the star formation (SF) history over the last  $\sim 0.15$  Gyr. The data are interpreted using theoretical simulations based on stellar evolutionary models. The synthetic diagrams include photometric errors and incompleteness factors. Testing various sets of tracks, we find that the ability of the models to reproduce the observed features in the CMD is strictly related to the shape of the blue loops of the sequences with masses around  $5 M_{\odot}$ . The field of NGC 1569 has experienced a global burst of star formation of duration  $\gtrsim 0.1$  Gyr, ending  $\sim 5$ – $10$  Myr ago. During the burst, the SF rate was approximately constant, and, if quiescent periods occurred, they lasted less than  $\sim 10$  Myr. The level of the SF rate was very high: for a single-slope initial mass function (IMF) ranging from  $0.1$  to  $120 M_{\odot}$  we find values of  $3$ ,  $1$ , and  $0.5 M_{\odot} \text{ yr}^{-1}$  for  $\alpha = 3$ ,  $2.6$ , and  $2.35$  (Salpeter), respectively. When scaled for the surveyed area, these rates are approximately 100 times larger than found in the most active dwarf irregulars in the Local Group. The data are consistent with a Salpeter IMF, though our best models indicate slightly steeper exponents. We discuss the implications of our results in the general context of the evolution of dwarf galaxies.

*Subject headings:* galaxies: evolution — galaxies: individual: NGC 1569 — galaxies: irregular — galaxies: starburst — galaxies: stellar content

## 1. Introduction

Understanding the status and evolution of dwarf galaxies is of crucial importance in current astrophysics and cosmology. They are important ingredients in common scenarios of galaxy formation, either as building blocks or as left-overs of the formation process (Silk 1987). In addition, dwarf galaxies forming at redshift around and below 1 have been suggested to be responsible for the excess of faint blue galaxies seen in deep photometric surveys (Babul & Rees 1992; Babul & Ferguson 1996).

Dwarf galaxies come in three main classes: (i) dwarf spheroidals, which contain population II stars and have low surface brightness (e.g., Wirth & Gallagher 1984); (ii) dwarf irregulars, with various levels of star-formation activity and a high gas mass (Hunter & Gallagher 1986); (iii) starbursting dwarfs, including blue compact dwarfs (Thuan 1991), H II galaxies (Terlevich et al. 1991), and blue amorphous galaxies (Sandage & Brucato 1979). The boundary between dwarf and giant galaxies is usually taken to be around  $M_B = -16$  mag (Tammann 1994).

An intriguing property of dwarf galaxies is the pronounced dichotomy between the two main classes of dwarf spheroidals and dwarf irregulars: most dwarfs are either gas-poor spheroids or gas-rich disk-like systems, with very few transition objects (van den Bergh 1977). The relationship between the two classes is still poorly understood: attempts to unify the complex zoo of dwarf galaxies have not been fully successful (Binggeli 1994). Some effort has been made to understand the interrelationship between individual classes in terms of an evolutionary sequence (e.g., Gallagher, Hunter, & Tutukov 1984). Starbursting dwarfs can play a key role: a powerful starburst might strip a gas-rich dwarf irregular of its gas and transform it into a gas-poor dwarf-elliptical (Kormendy 1985; Marlowe 1997).

Understanding the star-formation histories in starbursting dwarf galaxies is a prerequisite for understanding evolutionary connections between different object classes

(e.g., Gallagher 1996) since during cosmologically brief periods of time their star formation increases dramatically. This has profound implications for the interstellar medium. Stellar winds and supernovae inject energy and may cause a ‘blow-out’, initiating a galactic superwind (Heckman 1995). Numerous examples of galactic superwinds are known from observations (e.g., Heckman, Armus, & Miley 1990) and their cosmological implications have been discussed (e.g., De Young & Heckman 1994). However, attempts to relate the superwind properties and the star-formation histories in individual galaxies are quite rare and do not go beyond a qualitative level (Heckman et al. 1990; Leitherer, Robert, & Drissen 1992). Yet establishing such a relation is crucial for theoretical models of the superwind hydrodynamics (Suchkov et al. 1994; Tenorio-Tagle & Muñoz-Tuñón 1997).

NGC 1569 (= UGC 3056 = Arp 210 = VII Zw 16 = IRAS 4260+6444) has been indicated by Gallagher, Hunter, & Tutukov (1984) as an outstanding object in their sample of active star forming galaxies. Indeed, it is a prime candidate for a detailed study of the star-formation history in a starburst galaxy since at  $D = 2.2 \pm 0.6$  Mpc it is the closest starburst galaxy known (Israel 1988a). Adopting the literature reddening  $E(B - V) = 0.56$  (Israel 1988a), photometry of individual stars is feasible with HST down to  $M_{V,0} \simeq -1.5$ . This corresponds to stars with mass of about  $3 M_{\odot}$  in their core-helium burning phase, and consequently to a look-back time of  $\sim 0.4$  Gyr (e.g., Schaller et al. 1992).

The properties of NGC 1569 are typical of a dwarf galaxy. With a distance modulus of  $(m - M)_0 = 26.71$ , its total absolute B magnitude is  $M_{B,0} \sim -17$  (Israel 1988a), which is intermediate between the two Magellanic Clouds. Its total mass and hydrogen content are estimated  $M \simeq 3.3 \times 10^8 M_{\odot}$  and  $M_H \simeq 1.3 \times 10^8 M_{\odot}$ , respectively (Israel 1988a). Numerous studies of the chemical composition exist. The published range of oxygen abundances is very narrow:  $12 + \log(O/H) = 8.25$  (Hunter, Gallagher, & Rautenkranz 1982), 8.37 (Calzetti, Kinney, & Storchi-Bergmann 1994), 8.26 (Devost, Roy, & Drissen

1997), 8.29 (González-Delgado et al. 1997), and 8.19 (Kobulnicky & Skillman 1997; Martin 1997). Searches for chemical composition gradients were done by Devost et al. and Kobulnicky & Skillman. No significant evidence for chemical inhomogeneities was found. If  $[O/Fe] = 0.0$  is assumed, the average metallicity of NGC 1569 is  $Z \simeq 0.25 Z_{\odot}$ , with about 0.2 dex uncertainty. Thus, NGC 1569 is a gas-rich system with SMC-like composition in a relatively early stage of its chemical evolution.

The most detailed study of the warm ( $\sim 10^4$  K) gas of NGC 1569 was done by Waller (1991), who found evidence for a link between the star-formation history and the morphology and kinematics of extended gas structures. The same connection is suggested by hot ( $\sim 10^7$  K) gas observed with the ROSAT and ASCA satellites (Heckman et al. 1995; Della Ceca et al. 1996). The extended X-ray emission is consistent with a starburst driven galactic superwind which could in principle lead to a large-scale disruption of the interstellar medium. Evidence for large numbers of supernovae was also found from the non-thermal radio spectrum (Israel 1988b).

One of the most spectacular properties of NGC 1569 are its ‘super star clusters’. These high-density star clusters were first detected and discussed by Arp & Sandage (1985) and Melnick & Moles, & Terlevich (1985) and later studied by O’Connell, Gallagher, & Hunter (1994), Ho & Filippenko (1996), De Marchi et al. (1997), and González-Delgado et al. (1997). The super star clusters reflect a recent starburst in which at least  $10^5 M_{\odot}$  of gas was transformed into stars within about 1 pc. Super star clusters are similar to young Galactic globular clusters observed shortly after their formation (Meurer 1995).

Pre-Costar HST studies of the field population were done by O’Connell et al. (1994) and Vallenari & Bomans (1996) (hereinafter VB). According to the latter authors, NGC 1569 experienced a strong starburst until a few Myr ago, when its activity subsided. The repaired HST offers the possibility of significant improvement over these previous studies.

We have therefore embarked on an extensive spectroscopic and photometric HST study in an attempt to confront the observational data with the predictions of theoretical models, and derive detailed informations on the star formation (SF) history in NGC 1569. In a first paper (De Marchi et al. 1997) we discussed the photometry of the SSCs, while in this paper we present the results for the resolved stars in the same field, which are interpreted in terms of the recent SF history via theoretical simulations. The data and their analysis are described in Section 2. The derived color-magnitude diagram and luminosity function are in Section 3. Our theoretical models are described in Section 4. They are compared with the observations in Section 5. In Section 6 we discuss our results and their implications for the evolution of dwarf galaxies, and we present in Section 7 our general conclusions.

## 2. Data reduction and analysis

The observations of NGC 1569 described in this paper were obtained with WFPC2 on board the HST on 1996 January 10 (UT). NGC 1569 was centered in the field of view of the planetary camera (PC) yielding an effective plate scale of  $0''.046 \text{ pixel}^{-1}$  and a  $35'' \times 35''$  field of view. The plate scale and field of view of the adjacent wide field chips (WF) are  $0''.1 \text{ pixel}^{-1}$  and  $80'' \times 80''$ , respectively. Images were obtained in three broadband filters, F380W, F439W and F555W, whose photometric properties are similar — yet not identical — to the Johnson  $U, B, V$  filters, and are described in detail in the WFPC2 handbook (Biretta 1996). Since the photometric quality of the F380W images turned out to be poor, it was not possible to measure stars fainter than  $\sim 24$  mag with photometric errors smaller than  $\sim 0.2$  mag. We have then concentrated our analysis on the F439W and F555W data only, and used the F380W images to check for spurious detections (see below).

A set of 16 deep exposures was collected for each of these filters, all of duration 630 s in F439W and 450 s in F555W. Images were taken at 4 different positions, dithered by a

few pixel in order to improve the sampling of the final combined images, to recover faint stars with reliability, and to eliminate high frequency noise. The whole dataset has been processed using the standard STScI pipeline procedure. After correction for geometrical distortion, images taken with the same filter have been registered and combined to remove cosmic ray hits. Total exposure times of the final combined images are  $\sim 10,000$  s and  $\sim 7200$  s in F439W and F555W, respectively. For each filter, a set of short exposures was also obtained to provide photometric measurements of stars which would saturate in the deep images. Exposures of 20 s, 40 s, and 60 s duration were obtained for the F555W, F439W, and F380W filters, respectively, at two different pointings and are described in De Marchi et al. (1997).

Many bright star clusters and several thousand stars are clearly visible in the deep WF+PC images. The automated star detection routine DAOFIND was applied to the data, by setting the detection threshold at  $5\sigma$  above the local background level. We carefully examined by eye each individual object detected by DAOFIND and discarded saturated stars, a number of features (PSF tendrils, noise spikes, etc.) that DAOFIND had misinterpreted as stars, as well as a dozen extended objects (star clusters) whose full width at half maximum (FWHM) exceeded that of typical stellar objects in PC frames (FWHM  $\simeq 1.6$  pixel) by a factor of two or more.

In this way, a total of  $\sim 7000$  unsaturated stars were detected in the field covered by the PC. The severe crowding of our images, particularly towards the center of the galaxy, led us to prefer the use of the PSF-fitting DAOPHOT photometry package over the simpler, but more limited, aperture photometry techniques. A sample PSF was built by using three relatively isolated and moderately bright stars ( $m_{F439} \simeq m_{F555} \simeq 21.5$ ) located on opposite sides with respect to the body of the galaxy. Instrumental magnitudes measured in this way were converted into the WFPC2 synthetic system VEGAMAG as indicated in



Holtzman et al. (1995). Interested readers are referred to that paper for further details on the characteristics of the WFPC2 photometric system. In this paper,  $m_{F439}$  and  $m_{F555}$  refer to the WFPC2 synthetic system. Although these magnitudes are very similar to Johnson  $B$  and  $V$  magnitudes, no attempt has been made to convert our magnitudes into the standard Johnson system.

The main source of uncertainty in our photometry is crowding, particularly in the central regions of NGC 1569, where the average star-to-star distance is on the order of  $\sim 4$  pixels. The photometric error estimated by DAOPHOT ( $\sigma_{DAO}$ ), and confirmed by artificial star tests (see below), is smaller than 0.1 mag in both bands down to  $m_{F439} \simeq m_{F555} \simeq 24$ , and it grows to  $\sim 0.3$  mag at magnitude 26 in both bands. The behavior of the photometric error with magnitude is shown in Fig. 1.

We carried out a series of artificial star tests in order to estimate the fraction of stars lost to crowding as a function of their brightness. We ran 10 trials in each of the two filters, by adding each time a number of artificial stars not exceeding 10 % of the total number of objects in the frame, to avoid overcrowding. The magnitudes of the artificial stars were distributed according to the empirical luminosity function. Recursive DAOFIND runs, with the same parameters as in the reduction of the scientific data, allowed us to assess the fraction of added stars that were recovered and the magnitudes at which these objects were detected. Since the artificial star routine (ADDSTAR) places objects at random over the area of the frame, these tests provide an average value of the photometric incompleteness in our images. Artificial star tests were run on both filter frames, but the F439W image, being less deep, in practice defines the overall completeness limit. All artificial stars not recovered in the F555W frames were also missing in the F439W ones. This is partly due to the intrinsic color of the stars, and partly to the lower efficiency of the instrumental setup (WFPC2 + F439W) in the bluer band. Since the assumed color distribution of the

artificial stars follows the mean locus observed in the color-magnitude diagram (CMD), the incompleteness factors derived in the two filters are not independent and, as such, the overall incompleteness fraction is set by the larger of the two values at each magnitude.

For each artificially added star we compared the input magnitude with that at which the star was recovered, and found no systematic difference. In addition, the standard deviation of the recovered values is in excellent agreement with the  $1\sigma$  uncertainty of the photometric reduction at any magnitude level.

Our simulations show that at  $m_{F439} \simeq 25.0$  on average about 30 % of the artificial stars added to the PC chip are not detected by the automated routine, but this number goes down to only 8 % at  $m_{F439} \simeq 23.5$ . At magnitude levels brighter than  $m_{F555} \simeq 23$  all artificial stars are detected, except when they fall on top of one of the bright super star clusters. Our photometry is, therefore,  $\sim 100\%$  complete for  $m_{F555} < 23$ . Table 1 shows the completeness factors as a function of the F439W magnitude, as determined by the artificial star tests. These are the values adopted in our theoretical simulations.

The artificial star experiments provide us with a statistical estimate of the number of stars lost to crowding. There is, however, another issue that must be considered when comparing the observations with theoretical simulations: blending affects the derived magnitude of detected objects. The extent to which the images of two stars can merge with each other, thereby resembling a single object with photometric properties different from those of the parent stars, is a function of their angular separation and of their magnitude difference. If the projected angular distance between two stars is smaller than the PSF-FWHM, they will be detected as a single object by the photometric routine, and the number of stars lost because of this is included in the incompleteness factors. Concerning the photometric error, we select objects with  $\sigma_{DAO} < 0.1$  mag and take 0.1 mag as an estimate for the photometric accuracy of the data. Thus, we are essentially interested in

those cases in which the uncertainty introduced by blending on the determined magnitudes is larger than this value. This occurs when the blended stars differ in magnitude by less than  $\sim 2$  mag.

Our tests show, however, that even in this case the sharp PSF of the PC camera considerably limits the effects of blending, in spite of the high density of stellar objects in our frames. The results of the artificial star experiments, as well as statistical considerations, indicate that the probability of blending for a pair of stars differing less than 2 mag in flux amounts to  $\lesssim 1\%$  for objects brighter than  $m_{F555} \simeq 25$ . Therefore, it is unlikely that blending introduces a large photometric uncertainty. Blending of objects with a large magnitude difference is much more frequent. In this case, the fainter component is virtually lost with a small effect on the brighter one, and this is accounted for in the simulations by correcting for photometric error and incompleteness.

Two of the WF frames have also been reduced, the third having been discarded because of the presence of a highly saturated star. We found  $\sim 1000$  objects with  $\sigma_{DAO} < 0.2$  mag in both the F555W and F439W filters. Since the general characteristics of the stellar distributions in the CMDs derived for the WF and for the PC frames are very similar, we proceeded with a quantitative analysis only for the CMD derived from the PC frames.

### 3. Color-magnitude diagram and luminosity function

Fig. 2 shows the CMD in the HST F439W and F555W bands resulting from the analysis of the region of NGC 1569 contained in the PC field of view. As discussed by De Marchi et al. (1997), this region contains three super star clusters (SSCs), two of which form a close pair which is barely resolved. Only integrated photometry is feasible for the SSCs, which are not resolved into stars. The resulting  $(B - V)$  and  $(U - B)$  colors are

indicative of young ages, but is not possible to date the clusters precisely.

The diagram in panel a) shows the 2809 objects measured in both the F555W and F439W bands with a photometric error  $\sigma_{DAO} < 0.2$ , while the diagram in panel b) shows the subsample of 801 objects with  $\sigma_{DAO} < 0.1$ . By comparing the two panels it is apparent that the major features of the galaxy stellar population are well delineated in both cases. Clearly, the diagram with the tighter selection criterion contains a lower number of stars because it reaches a brighter magnitude limit, and because it rejects most of the objects with extreme colors.

In the CMD shown in panel a) there are some very blue objects, populating the bottom left corner. Taking into account the average extinction affecting our field, objects appearing with  $(m_{F439} - m_{F555}) \lesssim -0.5$  have intrinsic colors bluer than  $\simeq -1$ . However, neither galactic nor extragalactic objects can have colors that blue: both white dwarfs with magnitudes in the relevant range, and the bluest QSOs have intrinsic  $(B - V) \gtrsim -0.6$ . We have thus examined in detail all the 72 objects with  $(m_{F439} - m_{F555}) < -0.5$  and photometric error  $\sigma_{DAO} < 0.2$  mag in both bands. They are all concentrated in the region of the PC field dominated by the main body of NGC 1569, suggesting that these extremely blue objects originate from phenomena associated with the galaxy (either physical or spurious), and not from background or foreground effects. A detailed examination in all of the three photometric bands revealed that most of these objects are spurious detections, due to noise peaks falling on the wings of stellar images, or to the blend of 2 stars. When selecting only the objects with errors  $\sigma_{DAO} < 0.1$  mag no point source with extremely blue color is found. Since the shape and position of the bulk of the blue plume and of the red tail look unaffected by the selection criterion, we will adopt the diagram in panel b) as reference observational CMD for the stellar population in the field of NGC 1569.

The general morphology of the CMD of NGC 1569 is similar to that of Local Group

irregulars observed from the ground (see, e.g., Freedman 1988; Tosi et al. 1991, hereinafter TGMF; Greggio et al. 1993, hereinafter GMTF; Marconi et al. 1995, hereafter MTGF; Gallart, Aparicio, & Vílchez 1996b), with a quite scattered distribution and a prominent concentration of stars in the blue plume. The CMD shows a smaller fraction of red stars, but this is mostly due to the selection effect of the magnitude limit for object detection in the F439W frame. Indeed, the  $I$  vs.  $(V - I)$  diagram of NGC 1569 obtained VB shows a significant population of red (mostly intermediate age) stars.

The median color of the blue plume ( $m_{F439} - m_{F555}$ )  $\simeq 0.45$  is quite red as a consequence of high foreground extinction. By correcting this color for the average foreground reddening  $E(B - V) = 0.56 \pm 0.1$  as estimated from UV observations by Israel (1988a), an intrinsic  $(m_{F439} - m_{F555})_0 = -0.1$  is derived. This is equal to or slightly redder than the median  $(B - V)_0$  color of the blue plumes in other nearby irregulars (see, e.g., MTGF and Gallart et al. 1996b for NGC 6822; TGMF for Sextans B; Bresolin et al. 1993 and GMTF for NGC 3109). However, since González-Delgado et al. (1997) have derived  $E(B - V) = 0.67 \pm 0.02$  for the super star clusters in NGC 1569 (using UV, optical and near-IR spectra), it may well be that this galaxy is affected by internal differential reddening. Thus, the intrinsic colors of the blue plume stars could be bluer than the previous estimate.

Given the low galactic latitude  $b = +11^\circ$  of NGC 1569, the observed CMD is likely to be populated by a significant number of foreground stars. Despite our original request, the time allocated on HST was not sufficient to take exposures of adjacent fields and measure directly the Galaxy contamination. Since the CMD derived from the WF images looks very similar to that derived from the PC frames, the galaxy extends substantially over the WF frames, which cannot then be safely used to discriminate between members and foreground/background objects. However, using ground based observations, VB derive a very low density of external field objects in the direction of NGC 1569. Thus our CMD

should be almost unaffected by foreground/background contamination.

The full dots in Fig. 3 represent the luminosity function (LF) of the 801 selected stars of NGC 1569, whereas the open circles correspond to the stars with  $0 \leq (m_{F439} - m_{F555}) \leq 0.5$ , which in the following will be referred to as the *reference blue plume*. This subsample has been selected simply for the purpose of comparing our results with those relative to other galaxies and/or other authors, and does not represent either the whole blue plume or the pure main sequence (MS) stars.

In the past, the LF of the blue plume was usually adopted for estimating the mass function, under the assumption that it is essentially populated by MS stars. However, as already noticed for other irregulars (e.g., Freedman 1988; TGMF and GMTF; Gallart et al. 1996b), the comparison of the observational CMD with theoretical stellar models demonstrates that the blue plume is populated also by a significant fraction of stars at the hot edge of the blue loop evolutionary phase. This represents a serious problem for authors deriving the initial mass function (IMF) from the luminosity functions of the blue plume assuming it contained only MS stars. The MS–post-MS blend is apparent also in the  $I$  vs.  $(V - I)$  diagrams described by VB. As discussed there and in the next section, it is virtually impossible to discriminate MS from post-MS stars at the brightest magnitudes, where the morphology of the theoretical evolutionary tracks of massive stars in the CMD leads to the merging of the blue edges of the loops into the region populated by MS stars. Our  $(m_{F439} - m_{F555})$  selected *reference blue plume* is therefore just a compromise between the requirement of removing as much as possible the evolved stars and the need of keeping the brightest MS objects.

The slopes of the LFs are usually derived with least squares linear fitting to their complete portions. However, as shown in GMTF and MTGF, the large statistical fluctuations due to the small number of sampled stars, especially at the brightest

magnitudes, may significantly alter the results of a method requiring data binning. To overcome this problem, we have applied a maximum likelihood fitting to evaluate the slope of the LF in the *reference blue plume*. Down to  $m_{F555} = 23$ , where our sample is complete (see previous section), the slope turns out to be  $\Delta \log N/\Delta V = 0.66 \pm 0.05$ . This value is higher than  $0.46 \pm 0.06$  derived by VB, a result of the different methods: for instance, a linear fit gives a flatter slope of  $0.6 \pm 0.1$  for our data set. Furthermore, the fit of VB is not restricted to the data portion where completeness is reached.

Our slope  $\Delta \log N/\Delta V = 0.66 \pm 0.05$  is consistent with the average  $0.70 \pm 0.03$  derived by Freedman (1985) and Hoessel (1986) from a large sample of irregulars. It appears, however, slightly steeper than the slopes derived by us in a homogeneous way (see TGMF; GMTF; MTGF) for four nearby dwarf irregulars, whose average value is  $\Delta \log N/\Delta V \simeq 0.55$ .

#### 4. Theoretical models

The observed CMD and the (total) LF described in the previous section have been compared to corresponding theoretical simulations, based on stellar evolutionary tracks. This method has proven to be a powerful tool to investigate the star formation history of galaxies with young stellar populations (see Tosi 1994; Greggio 1994).

Synthetic CMDs are constructed via Monte Carlo extractions of (mass, age) pairs, according to a prescribed IMF and a star-formation rate (SFR) law. The extracted object is placed on the theoretical H–R diagram (HRD) by interpolating on evolutionary tracks, and its luminosity and effective temperature are converted into magnitudes using tables for bolometric corrections and colors. A photometric error and incompleteness test, both as measured on the observational frames, is finally applied, and the synthetic star is placed

on the CMD. Once the number of objects populating the synthetic CMD equals that of the observed one, the procedure is stopped, yielding the quantitative level of the SFR consistent with the observational data, for the prescribed IMF and shape of the SF law. This procedure automatically accounts for the stochastic nature of the SF process, small number statistics, photometric errors, and incompleteness factors affecting the real data. In addition, it easily allows us to explore the effects of using different sets of stellar tracks.

We have considered several sets of stellar tracks (Table 2) in order to compare their effect on the derived SF history, and to check their adequacy in reproducing the main features of the observed CMD. The interpolation algorithm was carefully chosen to reproduce the main characteristics of the evolutionary sequences. Due to numerical effects, the short lived, post-MS phases may be underrepresented in the Monte Carlo simulations. Thus, the tracks have been trimmed in subphases, and for each extracted object the evolutionary subphase is first determined. Interpolation between adjacent tracks is then performed within the appropriate subphase. This procedure ensures that the main features of the synthetic diagrams, (e.g., the blue-to-red ratio, the ratio between the MS and post-MS objects, the number of W-R stars, etc.) are consistent with the adopted set of tracks.

The luminosity and effective temperature of the synthetic objects are converted into magnitudes and colors by interpolating on tables constructed from the collection of model atmospheres by Lejeune, Cuisinier, & Buser (1997), which cover (although non-uniformly) the effective temperature and gravity ranges:  $2000 \text{ K} < T_{\text{eff}} < 50,000 \text{ K}$ ,  $-1.02 < \log g < 5.5$ , and are characterized by several values of the metallicity ( $[\text{Fe}/\text{H}]$ ), with solar elemental ratios. It should be noticed here that the tracks used for the simulations do not reach very low gravities, or temperatures lower than  $\sim 3600 \text{ K}$ . Therefore, the model atmospheres used for our synthetic diagrams are essentially those of Kurucz (1992). The sets at metallicities



$[\text{Fe}/\text{H}] = -0.5$  and  $[\text{Fe}/\text{H}] = -1$  have been selected for our application.

The conversion tables have been prepared as follows. For each model atmosphere in the grid, a theoretical bolometric correction  $BC_{V,th}$  has been computed:

$$BC_{V,th} = -2.5 \log \frac{\int_V L_\lambda t_\lambda d\lambda}{L_{bol}} \quad (1)$$

where  $t_\lambda$  is the  $V$ -filter transmission function in the VEGAMAG system. Similar tables for the  $m_{F555}-V$  and  $m_{F439}-V$  colors (in the VEGAMAG system) have been produced by processing the model atmospheres through the CALCPHOT routine in the SYNPHOT package. For the objects populating our synthetic HRD, the following relation applies

$$M_V = BC_{V,th} - 2.5 \log L_{bol} + C_V \quad (2)$$

where  $C_V$  is a calibration constant. In order to ensure that the theoretical magnitudes are in the *same photometric system as the observational CMD*,  $C_V$  has been fixed by enforcing that for the solar model atmosphere ( $T_{\text{eff}} = 5777$  K,  $\log g = 4.4$ ,  $[\text{Fe}/\text{H}] = 0.$ ), we obtain  $M_V = 4.7785$  if we adopt  $L_{bol,\odot} = 3.96 \times 10^{33}$  erg s $^{-1}$ . This is the value in the VEGAMAG system found by SYNPHOT when *observing* a star with the same flux distribution, the solar radius, at a distance of 10 pc. In this system, the solar model and the model atmosphere representing Vega ( $T_{\text{eff}} = 9400$  K,  $\log g = 3.9$ ,  $[\text{Fe}/\text{H}] = -0.5$ ) have  $(m_{F439} - m_{F555}) = 0.717$  and 0.0066, respectively.

For each synthetic object, characterized by (mass,  $\log L$ ,  $\log T_{\text{eff}}$ ), the theoretical bolometric correction and colors are determined through interpolation in these tables.  $M_V$  is then derived from eq. (2), and henceforth also the HST magnitudes in the VEGAMAG system.

## 5. Synthetic CM diagrams

The synthetic CM diagrams (CMD) and LFs depend on the input parameters in the simulation: IMF, history of the SFR, distance, reddening and theoretical stellar evolutionary tracks. Adopting a value for the distance modulus and reddening, the superposition of the evolutionary tracks to the observed CMD yields the range of ages and masses sampled by our data. As an example, in Fig. 4 we show the Geneva tracks with  $Z=0.001$  transformed to the observational plane, having adopted the canonical values  $(m - M)_0 = 26.71$ ,  $E(B - V) = 0.56$  from the literature. Comparing Fig. 4 with Fig. 2, panel a), one derives that the bluest stars in the blue plume are MS objects with masses larger than  $\sim 9 M_\odot$ , and then ages younger than  $\sim 30$  Myr. The wide color distribution shows the presence of evolved stars, with masses larger than  $\sim 4 M_\odot$ , and then ages younger than  $\sim 0.15$  Gyr. Thus, SF has been active in our field over  $\sim 0.15$  Gyr. Although the smoothness of the star distribution in the CMD qualitatively suggest a continuous kind of SF, this issue can only be quantitatively assessed computing the simulations, which take into account the smearing effect of photometric errors and the incompleteness at faint magnitudes. In our procedure, then, we first try the simplest possible case, i.e. a constant SF over the look-back time, and then vary the SFR shape and IMF slope to reproduce the data distributions. When an acceptable representation of the observation is reached, we perturb the parameters to test the stability of the solution.

The comparison of the theoretical simulations with the observations is performed quantitatively on the LF, and mostly qualitatively on the appearance of the CMD. A detailed quantitative comparison of the color distribution of stars (like e.g., Tolstoy & Saha 1996) would not add much information, due to the smearing effect of photometric errors, and, more importantly, to the intrinsic uncertainty affecting the effective temperatures of models of massive stars in the post-MS phases (Renzini et al. 1992). Small variations of the

theoretical  $T_{\text{eff}}$  result in large differences of the predicted distribution of objects throughout the CMD, due to the high sensitivity of colors and bolometric corrections to the effective temperature in the relevant range.

Because of systematic uncertainties, there is often no significant difference between the best simulation and other similar solutions. Rather, our aim is to isolate the range of IMF and SFR parameters which are consistent with the data. We have computed a few hundred synthetic CMDs varying the input parameters mentioned above. In the next subsections we discuss our results and the influence of the input parameters on the conclusions. We start with adopting a distance of 2.2 Mpc, and present our preferred solutions for various sets of evolutionary models. Second we discuss how different SF histories influence the results. Finally we consider the effect of an uncertainty in the distance. The reddening is varied, within the observational estimate, in such a way that the average color of the synthetic blue plume agrees with the observations.

### 5.1. Influence of the Evolutionary Sequences.

The SF history derived from the analysis of an observational CMD depends on the set of tracks adopted in the simulation (Greggio 1994). We have therefore considered several sets of stellar tracks (Table 2). Although the metallicity of NGC 1569 is  $Z \sim 0.004$  we have computed simulations based on the Geneva tracks with several values of  $Z$ . This is motivated by the fact that the overall appearance of the CMD is sensitive to the morphology of the blue loops, which in turn critically depends on  $Z$  (Ritossa 1996).

We have also explored the difference in the derived interpretation when adopting the Padova set of tracks at  $Z = 0.004$ . The main difference between the Padova and the Geneva sequences concerns the treatment of convective overshooting: both sets include core

overshoot, but to a lesser extent in the Geneva models. In addition, the Padova tracks also include overshooting from the base of the convective envelope.

It is worth recalling some general features of the evolutionary sequences:

- i) All sets are characterized by extended loops for masses above  $4 M_{\odot}$  with the maximum extension for the Geneva  $Z = 0.001$  and Padova  $Z = 0.004$  sets.
- ii) Among the Geneva tracks, the fraction of evolutionary lifetime spent in the blue portion of the CMD during the post-MS phase is largest for the  $Z = 0.001$  tracks (actually close to 100% for stars more massive than  $\sim 4 M_{\odot}$ ), and it decreases with increasing metallicity. The Padova set at  $Z = 0.004$  is characterized by a value of this ratio which is intermediate between the Geneva  $Z = 0.001$  and  $Z = 0.004$  sets.
- iii) The ratio between post-MS and MS evolutionary lifetimes for stars more massive than  $\sim 8 M_{\odot}$  is  $\sim 0.1$  for the Geneva tracks. The same ratio is considerably smaller for the Padova tracks, reaching a minimum value of  $\sim 0.06$  for the  $12 M_{\odot}$  track.

These characteristics crucially determine the main differences in the simulations constructed with the various sets of tracks.

#### 5.1.1. *Simulations with the Geneva tracks with $Z = 0.008, 0.004$*

Adopting the Geneva tracks with  $Z = 0.008$  and  $Z = 0.004$  we were not able to reach a satisfactory agreement between the synthetic diagrams and the observations, regardless of the choice of the IMF and SF parameters. As an example, in Fig. 5 we show two of the best cases obtained with  $Z = 0.008$  (panel a) and  $Z = 0.004$  (panel b). The corresponding LFs are shown in panel c), with the solid line referring to the simulation displayed in panel a). The adopted reddening values are  $E(B - V) = 0.46$  and  $0.56$  for panels a) and b)

respectively, which compensate for the different metallicities of the tracks, and are both consistent with Israel’s (1988a) estimate.

The CMD in panel a) has been computed with two partially overlapping SF episodes, so that the SFR peaks around  $\sim 0.05$  Gyr ago. The adopted IMF exponent is  $\alpha = 3$  (in these units the Salpeter’s exponent is 2.35). The simulation in panel b) adopts a mildly decreasing SFR, and a Salpeter IMF. In both simulations the synthetic blue plume extends up to too bright magnitudes and the bright portion of the LF is overpopulated over a wide magnitude range. These characteristics appear in all the simulations with this sets because they are related to the shape of the tracks on the CMD. Infact, due to the relatively short color extension of the blue loops of models with mass around  $5 M_{\odot}$  (see Fig. 6 for the  $Z = 0.008$  set), the faint portion of the blue plume has to be mainly populated by MS stars. This is a consequence of the photometric incompleteness in the F439W frames, which depletes the number of evolved stars when their colors are too red. Thus, with these tracks a large number of massive stars is required, whose post-MS progeny shows up as bright evolved supergiants. In the specific case of the  $Z = 0.008$  tracks, at magnitudes brighter than  $m_{F555} \simeq 21.5$  we find post-MS objects with masses between 15 and  $25 M_{\odot}$ , which are the evolved progeny of fainter MS objects populating the blue plume in the range  $m_{F555} \simeq 25 - 22.5$ . Thus, the excess of bright objects cannot be suppressed without substantially underpopulating the faint portion of the blue plume, and could only be avoided by invoking such a delicate fine-tuning of the SF behavior, to appear very contrived. Similar arguments apply to the  $Z = 0.004$  set, in which the extension of the blue loops of the  $5 M_{\odot}$  and  $7 M_{\odot}$  tracks is so short, that the Hertzsprung gap appears at the faint magnitudes, in spite of the large photometric error.

The CMDs in Fig. 5 were computed with the low mass loss option in the Geneva models. Switching to the high mass-loss sets does not improve the comparison between the

simulations and the data. For  $Z = 0.008$ , the higher mass loss only affects the location of the core helium burning phase for the more massive models. As a consequence, the excess objects are still present, yet shifted to intermediate and red colors. For  $Z = 0.004$ , the adoption of a high mass loss rate would only affect the evolution of the most massive stars ( $M \gtrsim 40 M_{\odot}$ ), spanning a mass range which is almost not populated in our CMD.

### 5.1.2. Simulations with the Geneva tracks with $Z = 0.001$

As can be seen in Fig. 4, the loops for the Geneva tracks with  $Z = 0.001$  extend to very blue colors, and a substantial fraction of the faint blue plume can be populated by relatively low mass stars in their core helium burning phase. Unlike the previous cases, we find satisfactory agreement with the observed CMD and LF with various combinations of the IMF slope and SFR, provided that the activity ceased  $\sim 7$  Myr ago (corresponding to the evolutionary lifetime of a  $\simeq 30 M_{\odot}$  star) to avoid an excess of bright stars. Fig. 7 shows our best models for a constant SFR (panel a) and an exponentially declining SFR with an e-folding time of 0.6 Gyr (panel b). Notice that this latter case implies a variation of the SFR of only  $\sim 20\%$  over our short look-back time. The corresponding LFs are displayed in panel c). A reddening of  $E(B-V)=0.56$  has been adopted for these simulations. It can be seen that both cases give a very good fit to the observed LF. We adopted a steeper IMF for the constant SF case, in order to avoid too many bright stars. The synthetic CMDs also give a fair representation of the data, except for a slight paucity of stars at intermediate and red colors.

In order to see whether the data allow for large variations of the SF activity during the last 0.15 Gyr we have computed simulations with two separate SF episodes. Fig. 8 shows our best models obtained by assuming that the SFR in the most recent episode is higher (panel a), and lower (panel b) with respect to the previous episode. In both cases the SFRs

in the two episodes differ by a factor of  $\sim 1.2$ . Again, we achieve a fair representation of the data when appropriately tuning the IMF slope with the SF history. In all acceptable simulations, however, the SFR in the last 0.15 Gyr has proceeded approximately at the same level, and the slope of the IMF is close to Salpeter’s value or slightly steeper.

Our procedure corresponds to counting and weighing the stars which appear in the observational CMD, and determine their ages. The derived level of the SFR is then relatively insensitive to the combination of IMF and SF history adopted in the simulations. Indeed, our models in Fig. 7 and Fig. 8 show agreement with the observations, in spite of the different prescriptions. Yet, the average values of the SFRs over the last 0.15 Gyr, in stars more massive than  $4 M_{\odot}$ , are  $0.07$  and  $0.09 M_{\odot} \text{ yr}^{-1}$  for the simulations in Fig. 7 (panels a and b respectively), and  $0.08$  and  $0.11$  for the simulations in Fig. 8 (panels a and b respectively). This applies only to the mass range directly observed. When extrapolating the SFR for stars with masses ranging from  $0.1$  to  $120 M_{\odot}$ , the average rates become  $\sim 3 M_{\odot} \text{ yr}^{-1}$  for  $\alpha = 3$  and  $\sim 0.5 M_{\odot} \text{ yr}^{-1}$ , for  $\alpha = 2.35$ .

### 5.1.3. Simulations with the Padova tracks with $Z = 0.004$

Unlike the Geneva tracks with the same metallicity, the Padova tracks with  $Z = 0.004$  are characterized by well extended loops for masses larger than  $\sim 5 M_{\odot}$  whose blue edges almost merge into the MS. As a consequence, the faint blue plume is populated by MS and helium burning stars. Moreover, in the range  $9 - 25 M_{\odot}$  the ratio between post-MS and MS lifetime is significantly smaller than in the Geneva tracks, implying a lower ratio between bright evolved supergiants and their MS progenitors. For these reasons the synthetic diagrams with this set of tracks are fairly different from those obtained with the Geneva tracks with the same metallicity. We are able to obtain a fair representation of the data, provided that the activity stopped  $\sim 8 - 10$  Myr ago to avoid an excess of bright stars.

By adopting the spectroscopic value for the reddening,  $E(B - V)=0.56$ , the color of the synthetic blue plume appears somewhat redder than the observed one. We have therefore computed cases with both  $E(B - V) = 0.56$  and 0.46.

Although assuming a constant SFR we obtain acceptable simulations, the best fitting models are characterized by two distinct episodes of SF, the most recent one starting  $\sim 30$  Myr ago. We find acceptable solutions with SFRs both increasing and decreasing with time, in combination with different values of the IMF slope and adopted reddening. However, all the solutions indicate that the SFR has not varied by a large amount within the last 0.15 Gyr, and that the IMF slopes are slightly steeper than Salpeter’s value, and close to that derived for solar neighborhood stars (e.g., Scalo 1986). As an example, Fig. 9 shows the synthetic CMDs obtained with two bursts, the most recent one occurring at a lower rate in both panels a) and b). The theoretical and observed CMDs and LFs are in good agreement, perhaps with a slight excess of red supergiants in the models. In these models the ratio between the SFR in the older and most recent episode is 1.5 (panel a) and 1.2 (panel b). Thus, also with this set of tracks the data are consistent with an almost constant SFR over the last 0.15 Gyr (within 20% to 50%). The average level of the SFR in the simulations in Fig. 9 are 0.1, 0.07  $M_{\odot} \text{ yr}^{-1}$  in panels a and b respectively, in stars younger than 0.15 Gyr and more massive than 4  $M_{\odot}$ . When extrapolating the IMF down to a lower mass cut-off of 0.1  $M_{\odot}$  we obtain an average SFR of is  $\sim 1 M_{\odot} \text{ yr}^{-1}$  if  $\alpha = 2.6$ , or  $\sim 3 M_{\odot} \text{ yr}^{-1}$  if  $\alpha = 3$ . This holds when adopting  $E(B - V) = 0.46$ . For the higher reddening the derived rates are  $\sim 1.3$  times larger.

## 5.2. Testing the SF history.

Given the smooth distribution of stars in the CMD, and the shape of the LF, large variations of the average SFR within the last  $\sim 0.15$  Gyr are not consistent with the data,



as we have tested numerically. In this section we rather concentrate on the epoch of the recent stop in the SF, and on the possibility of short quiescence periods.

Fig. 10 shows the sensitivity of the results on the adopted age for the recent stop in the SF in the case of the Geneva tracks with  $Z = 0.001$ . In the bottom panel the SF has ceased 5 Myr ago, while in the top panel it is still ongoing. A steep IMF has been adopted for these simulations, in order to minimize the number of massive stars. Despite that, an ongoing SF predicts an excess of bright objects, while a 5 Myr stop is consistent with the data. Fig. 11 panel b) shows the case of a constant and currently ongoing SF having adopted the Padova  $Z = 0.004$  models. In spite of the steep exponent of the IMF, the CMD turns out to be overpopulated at the bright end, over a range of  $\sim 1.5$  mag. With this set of tracks, adopting  $\alpha=3$ , we still find an excess of bright stars stopping the SF 7 Myr ago. It is worth noticing that the age of the stop is constrained by the bright portion of the LF, over a range of  $\sim 1$  mag or more.

We now turn to consider the possibility of discrete episodes of SF, separated by quiescent phases. In general, this behavior has a definite signature on the CMD, i.e. the lack of objects around the isochrones with ages corresponding to the quiescence period, which shows up in the evolved portion of the diagram as an almost horizontal gap. Due to the effect of photometric errors, however, this signature can go undetected on the observed CMD.

A quiescent period lasting from 35 to 30 Myr ago has been assumed in the simulations shown in Fig. 9, in order to improve the appearance of the synthetic CMDs. The simulations with a constant SFR are characterized by a clump of blue supergiants at  $m_{F555} \simeq 22$  (see Fig. 11, panel b) populated by objects with masses in the range  $\sim 9$  to  $12 M_{\odot}$ . The blue loops of these sequences happen to converge to the same location in the CMD, providing an excess of objects in this region. Isochrones with ages of 35 to 30 Myr are populated by

post-MS objects with masses in the range  $\sim 9$  to  $10 M_{\odot}$  and the assumed quiescent period makes this clump of blue SGs less prominent. It should be noticed that a slightly different shape of the loops of these sequences would not produce this feature, making the quiescent period unnecessary. We also find that quiescent periods longer than  $\sim 10$  Myr should be excluded, since they show up in the synthetic CMD with a very evident horizontal gap, which is absolutely absent in the observed diagram. This is illustrated in the top panel of Fig. 11, where we show a simulation in which no SF has occurred between 40 and 30 Myr ago.

The simulations in Fig. 8, based on the Geneva tracks with  $Z = 0.001$ , have been computed assuming two contiguous SF episodes. Also for these tracks we have tested the possibility of short interburst periods. Again, we find that quiescent periods longer than  $\sim 10$  Myr in the recent past (less than 0.1 Gyr ago) would be evident in the observational CMD, and are then excluded by the smoothness of the data.

### 5.3. Dependence on the adopted distance.

The distance to NGC 1569 is relatively uncertain. The most widely quoted values in the literature are between 2.2 and 2.5 Mpc (Arp & Sandage 1985; Israel 1988a; O’Connell et al.1994). They are based on the properties of the still resolved brightest individual supergiants. However, distances of up to 4 Mpc are possible since NGC 1569 is a probable member of the IC 341/Maffei 1 & 2 group, whose distance is uncertain. If NGC 1569 were at 4 Mpc, it would be quite extraordinary in several respects. Apart from overluminous individual supergiants it would harbor the most luminous resolved super star clusters known in the universe. The brightest cluster has  $M_V = -13$  to  $-14$  already for the shorter distance (O’Connell et al.1994; De Marchi et al.1997). At 4 Mpc, this cluster would have appeared at  $M_V \simeq -17$  if observed at peak brightness 8 Myr ago! A larger distance would

make the properties of the ring nebula in the outskirts quite unusual as well. Drissen & Roy (1994) noted that its size and  $H_\alpha$  luminosity exceeds those of similar ring nebulae in M33 by up to an order of magnitude. This apparent discrepancy would be even more amplified for the larger distance.

While none of these arguments by itself provides convincing evidence for or against one or the other distance, taken together they all point toward the shorter distance. However, a larger distance would imply younger ages for the stars in our observed CMD, as well as larger levels of the recent SFR. To test the effect of the uncertainty in the distance to NGC 1569 we have computed simulations assuming a distance of 4 Mpc. The observed CMD now samples stars younger than  $\sim 50$  Myr. Two of our best cases are shown in Fig. 12, for the Padova tracks with  $Z = 0.004$ . The simulations assume a still ongoing SF (bottom panel) and a recent stop, at 5 Myr (top panel), and again the slope of the IMF has been chosen to minimize the number of massive stars. It can be noticed that, even adopting a large distance, a stop in the recent SF activity is preferred, not to overpopulate the bright end of the LF. The level of the SFR in these simulations is higher by a factor of  $\sim 2.8$  with respect to what derived with the shorter distance.

## 6. Discussion

### 6.1. Summary of the main results

In the previous section we have shown that the Geneva tracks with metallicities  $Z = 0.008$  and  $0.004$  do not allow us to reproduce the main features of the observed CMD and LF, regardless of the adopted combination of IMF and SFR. On the contrary, we find a fair agreement between the simulations and the data when adopting either the Geneva tracks with  $Z = 0.001$ , or the Padova tracks with  $Z = 0.004$ , the latter value being

consistent with the metallicity derived observationally. These results, however, are mostly based on the shape of the blue loops, which are intrinsically uncertain. Therefore, we can neither conclude that there is an indication for a lower metallicity in this galaxy, nor that the input parameters used in the Padova tracks are to be preferred to those in the Geneva tracks, as our analysis does not allow us to constrain the shape of the tracks down to this level of detail. Rather, we maintain that both sets of tracks have the combination of input physics and metallicity parameter that yields a good representation of our data. The color distribution of the evolved stars cannot be used to determine the metallicity of the system, other than to derive a general ranking.

We can derive several consistent indications on the SF history in the field of NGC 1569, regardless the adopted sequences:

- i) The SF activity stopped 5–10 Myr ago.
- ii) During the last 0.1–0.15 Gyr the SFR has been approximately constant. If quiescent periods have occurred, they did not last longer than  $\sim 10$  Myr.
- iii) The best agreement is generally reached with IMF slopes steeper than Salpeter’s, but the latter would not be inconsistent with the data.
- iv) The average SFR in our field is approximately 3, 1, and  $0.5 \text{ M}_{\odot} \text{ yr}^{-1}$  for IMF slopes of 3, 2.6, and 2.35, respectively. These values refer to stars with masses ranging from 0.1 to  $120 \text{ M}_{\odot}$  but it should be emphasized that our data only constrain the star formation of objects more massive than  $\sim 4.5 \text{ M}_{\odot}$ . The dependence of these derived values on the adopted set of tracks, reddening and shape of the SF law is found to be small.
- v) Assuming that the minimum mass for a type II SN explosion is  $8 \text{ M}_{\odot}$  the derived SFR implies that in the last 0.1 Gyr in our region about  $2 - 3 \times 10^5$  type II events have occurred, for an IMF slope ranging from 3 to 2.35.

It may seem that enhancing the recent SFR while steepening the IMF would lead to the same result, and to some extent this is true: enhancing the recent SF would populate the CMD with young, and therefore massive, stars, thereby counterbalancing the effect of a steepening of the IMF (see Fig. 7 and Fig. 8). However, we find that acceptable solutions have parameters in a relatively small range. In other words, when the shape of the SFR is radically different from the one giving a fair solution, it is difficult to obtain acceptable simulations by varying the IMF slope (and vice versa). Indeed, the two parameters do not act in precisely the same way, as strengthening the recent SF with respect to the past enhances the population along the younger isochrones by a constant factor. Conversely, varying the slope of the IMF affects the distribution of stars along all the isochrones. Thus, for example, when the IMF slope is too flat, the LF of the synthetic population is so skewed towards the bright end that only very complicated SF histories may fit the data. As can be seen from Figs. 6 and 4, the shape of the tracks, and therefore of the isochrones as well, is such that in a given magnitude bin we find stars of different generations and different masses, so that it is not possible to disentangle the effects of SFR and IMF. The number of stars falling within suitably chosen boxes in the CMD (e.g., Bertelli et al. 1992) may allow this task. However, it is very hard to define these suitable boxes in this case, due to the shape of the tracks for massive models in the observed CMD and to the photometric errors. This cautionary note applies in particular to the splitting of the population of the blue plume into the MS and post-MS contributions. For these reasons we need theoretical simulations, rather than isochrone superposition, to interpret the data. Moreover, as shown by our experiments, the shape of the blue loops heavily affects the synthetic CMD and LF so that the final solution depends on the adopted set of tracks. The flexibility of the simulation code allows us to easily explore this parameter space as well.

It is also worth noting that the SF stop that we infer in the field of NGC 1569 is not particularly sensitive to the adopted upper mass cut off for the IMF. For example, when the

mass distribution is truncated at  $\sim 40 M_{\odot}$  the synthetic diagrams obtained with an ongoing SF are still overpopulated at the brightest magnitudes. Indeed, lowering the mass cut off does not appreciably alter neither the slope of the LF, nor the overall ratio between the objects populating the fainter and brighter portions of the LF. Conversely, when stopping the SF activity a few Myr ago, the whole MS population is reduced with respect to the evolved one.

Our quoted values for the SFR refer to a single-slope IMF with stars distributed from 0.1 to  $120 M_{\odot}$ , and their dependence on the lower mass cut-off is of course very high. For example, the mass in stars within 0.1 and  $1 M_{\odot}$  is  $\sim 0.55$ ,  $0.75$ , and  $0.9$  of the total for  $\alpha = 2.35$ ,  $2.6$ , and  $3$ , respectively. Furthermore, these values depend on the IMF slope at the low mass end. In a recent study by Gould, Bahcall, & Flynn (1997), the IMF for M dwarfs in the galactic disk can be fitted by a two-slopes IMF, with a break at  $M \simeq 0.6 M_{\odot}$  and with a slope  $\alpha = -0.56$  (in our usual units) in the low mass range. When adopting these prescriptions for the low mass end of the IMF, and varying the slope of the mass distribution only over the range  $M \geq 0.6 M_{\odot}$ , our average SFRs become  $\sim 0.8$ ,  $0.5$ , and  $0.3 M_{\odot} \text{ yr}^{-1}$  for slopes of  $3$ ,  $2.6$ , and  $2.35$ , respectively.

## 6.2. Comparison with previous work

The only previous study primarily dedicated to the resolved stellar population in the field of NGC 1569 is that by VB. By using pre-refurbishment WFPC images, these authors derive the CMD in the F555W and F785LP filters, roughly equivalent to the  $V$  and  $I$  bands. The method used for the interpretation of the data relies, like ours, on simulations of CMDs based on evolutionary tracks, but without including the effect of photometric errors and incompleteness. An additional difference concerns the use of the LF as a tool to discriminate between possible models.

VB conclude that NGC 1569 experienced a global burst of SF from 0.1 Gyr to 4 Myr ago, and that a previous episode of SF had taken place in the galaxy, at a substantially lower rate. Our observations are not deep enough in the F439W band to detect the AGB stars that reveal this older episode in the  $I$  vs  $(V - I)$  diagram. As for the recent SF history, our results are roughly consistent with their finding. We prefer an earlier stop of the SF activity, since, when adopting an SF rate ongoing up to 4 Myr ago, the synthetic LF tends to be overpopulated at the bright end, especially for the Padova tracks with  $Z=0.004$ , which is the same set adopted by VB. VB could not establish quantitatively this effect without the comparison of the theoretical and observed LFs. In addition, we believe that the photometric incompleteness prevents the determination of the epoch at which the burst started, since stars older than 0.15–0.1 Gyr would go mostly or totally undetected in our images. Finally, the level of the SFR of the recent burst that we derive using the same set of tracks and IMF is consistent with theirs, judging from the estimate of the recent SNII activity.

To summarize, our results are in general agreement with VB, but we believe that our conclusions are more robust, due to both the superior quality of the refurbished HST data, and to the method employed for the interpretation.

### 6.3. NGC 1569 as a post-starburst galaxy

NGC 1569 has been suggested to be in a post-starburst phase (Israel 1988a). The definition of the starburst and the post-starburst phase is a matter of choice. Here we define the starburst phase as being characterized by ongoing SF. With this definition the field population of NGC 1569 clearly is in a post-starburst phase since we find that the CMD requires no SF activity over the last few Myr.

On the other hand, in the same region there are many HII regions and three SSCs, whose age is likely to be younger than 10 Myr (González-Delgado et al. 1997). Therefore the stellar population in NGC 1569 is made of two components: the resolved stars with ages older than  $\sim 10$  Myr, and the stars in the SSCs, which formed more recently. The resolved stellar component is in a post-starburst stage, while the most recent star formation is taking place in the SSCs. This finding may suggest a picture of discrete SF events mainly generating SSCs, which dissolve to create the field stellar population in a certain time interval. The  $\sim 10$  Myr quiescence period would correspond to a characteristic time step of the discrete SF activity. On the other hand, the compactness of the SSCs suggests that they are gravitationally bound systems, and that they are true young globular clusters (Ho & Filippenko 1996). In addition, the age of the brightest component of SSC-A has been estimated to be as old as 8 Myr (González-Delgado et al. 1997), and still presents a half-light radius of  $\sim 1.6$  pc (De Marchi et al. 1997). Therefore the field population is likely to be the relic of a major episode of SF which involved the whole observed region, and which ended several Myr ago. The SF activity in the field is most likely responsible for the outflow of the interstellar medium analyzed by Waller (1991), who derived a kinematic age of 10 to 60 Myr for the H $\alpha$  filaments around NGC 1569. This SF activity, however, has not depleted the gas reservoir, and more recent SF is currently occurring in the HII regions and in the SSCs. This may suggest that only the densest regions are able to collapse to form stars following the burst of SF and its consequent energy input in the ISM.

#### 6.4. Comparison with other dwarf irregular galaxies

The SFR in dwarf irregulars has been found to be characterized by long activity periods, separated by short interburst phases (*gaspings SF*, e.g., Tosi 1994). In this respect, the field stellar population in NGC 1569 exhibits similar properties, with a SF occurring



at an approximately constant level over  $\sim 0.15$  Gyr, with quiescence periods shorter than  $\sim 10$  Myr. A quantitative comparison is instructive. Among the different possible ways in which the strength of the star formation activity can be characterized, we consider the SFR per unit surface (see Hunter & Gallagher 1986).

At a distance of 2.2 Mpc, the observed PC field surveys an area of approximately  $0.14 \text{ kpc}^2$ , so that our derived recent SF rates are about 20, 7, and  $4 \text{ M}_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$ , for IMF slopes of 3, 2.6, and 2.35, respectively (again having extrapolated the IMF down to  $0.1 \text{ M}_\odot$ ). It is worth recalling that the typical SFR quoted for the solar neighborhood (e.g. Timmes, Woosley, & Weaver 1995) is  $\sim 1000$  times lower than this value.

The comparison of the SFR in NGC 1569 with those derived for other dwarf irregulars is not straightforward since there is a non-trivial dependence on the method used to measure the SFR. Even when considering the SFR derived from the analysis of the CMD via theoretical simulations, there may be systematic differences due to the adopted set of stellar tracks (Greggio 1994). Finally, the SFR depends on the adopted IMF, both on the slope and on the mass cut-offs, the latter being not always clearly specified in the literature.

With this caveat in mind, the recent SFR derived via theoretical simulations for dwarf irregular galaxies in the Local Group range from approximately  $10^{-4} \text{ M}_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$  to  $10^{-2} \text{ M}_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$  (MTGF; Tolstoy 1996; Gallart et al. 1996a). When allowing for the uncertainties introduced by the different methods used, we can conclude that the recent SFR in NGC 1569 is  $\sim 100$  times larger than in the most active local dwarf irregular. NGC 1569 fully deserves its classification as a (post)-starburst galaxy, as opposed to normal, star-forming galaxies.

Could it be that strong SF bursts took place in dwarf irregulars in the past? The chemical properties, and the gas content of the most active of them are not consistent with the extrapolation of the currently observed activity over a long time. Chemical evolution

models show that several bursts of SF may have occurred in these galaxies, but separated by long quiescence periods. Strong bursts of SF could perhaps be accommodated but the duty cycle of this activity has to be low (Matteucci & Tosi 1985; Pilyugin 1994; Marconi, Matteucci, & Tosi 1994).

More direct information on the past SF activity occurred in dwarf irregulars has been derived for some Local Group objects by analyzing the stellar population along the giant branches (Gallart et al. 1996a; Aparicio, Gallart, & Bertelli 1997a,b). Although the SFRs derived in this way are less detailed and reliable than those deduced from the analysis of the MS and core He-burning stars, there is no hint that these galaxies experienced a much stronger SFR in the past than recently. On the other hand, the analysis of the giant branches does not permit a fair time resolution, and the past activity in these galaxies could also have been characterized by recurrent strong bursts, whose average rate is reflected by the bright portion of the CMD.

The CMD derived by VB exhibits a well populated AGB, which definitely argues for previous SF in NGC 1569. VB concluded that a SF episode took place at a significantly lower rate than the most recent one between 1.5 and 0.15 Gyr ago. However, the values of these parameters are uncertain due to observational and theoretical limitations. Nevertheless, the mere presence of AGB stars gives evidence for a previous SF activity at intermediate ages. Ages as old as  $\sim 10$  Gyr can be probed only with NICMOS observations looking for the presence of stars at the tip of the first red giant branch in the infrared.

In the meantime, indirect information on the past SF history can be derived from chemical evolution arguments. A recent determination of the oxygen, nitrogen and helium abundances in the interstellar medium of NGC 1569 was done by Kobulnicky & Skillman (1997). Comparing their values to the predictions of chemical evolution models for bursting dwarfs (Marconi, Matteucci, & Tosi 1994) we find that several bursts of SF may have

happened in this galaxy, provided that strong galactic winds have occurred. The products of SNII explosions should be preferentially lost in these winds, thus keeping the overall metallicity low.

To summarize, the comparison of the SF history in NGC 1569 and that of dwarf irregular galaxies in the Local Group is inconclusive. While the general characteristics of the SF appear very similar, the level of the current SFR in NGC 1569 seems significantly larger than that of the dwarf irregulars in the Local Group over their whole lifetime. This suggests that NGC 1569 is distinctively more active than the Local Group dwarf irregulars.

An evolutionary link between dwarf spheroidals and dwarf irregulars has been suggested several times (e.g., Kormendy 1985; Binggeli 1994). Strong bursts of SF are known to have occurred in dwarf spheroidals in the past (e.g., Smecker-Hane et al. 1994). Unfortunately, to our knowledge a quantitative determination of the level and duration of these bursts has not yet been attempted. The evidence that NGC 1569 is currently experiencing a galactic outflow seems well documented, an effect which would turn the galaxy into an inconspicuous, perhaps forever fading object. On the other hand, the ultimate fate of the expanding gas is rather uncertain, due to the complications of hydrodynamic modelling. A stronger case could perhaps be made by comparing the SFR in the bursts which occurred in the dwarf spheroidals, to what we measure for NGC 1569.

### 6.5. Implications for the faint blue galaxies

We now consider the possible relevance of galaxies like NGC 1569 to the interpretation of faint galaxy counts. To do this, we first discuss the possible evolutionary path for the SFR in this galaxy.

The direct census of the stellar population in NGC 1569 suggests that a strong burst

of SF has occurred at a rate of  $\sim 0.5 \text{ M}_{\odot} \text{ yr}^{-1}$ , with a duration of  $\sim 0.15 \text{ Gyr}$ . This level of the SFR cannot have been sustained over a long period of time. The total mass of the galaxy as estimated by Israel (1988a) is  $\sim 3.3 \times 10^8 \text{ M}_{\odot}$ . Considering a SFR of  $0.5 \text{ M}_{\odot} \text{ yr}^{-1}$  as representative of the whole galaxy (i.e. neglecting the activity in the regions external to our PC field, and considering a turn-over in the IMF at the low mass end) it only takes  $\sim 0.7 \text{ Gyr}$  to cycle the entire mass into stars. The gas exhaustion timescale is longer than this because of the mass return from stellar winds and supernovae. However, this corresponds to a small correction since 1 Gyr after its birth a single age stellar population has returned only  $\sim 0.2$  of its total mass (for a Salpeter IMF). Thus it seems very unlikely that such a high SFR could have been supported throughout a lifetime of several Gyr, even allowing for the occurrence of galactic winds in the past, which would imply a larger initial mass. Furthermore, the metallicity resulting from such extreme activity would inevitably be much larger than derived from the spectra of HII regions. Though the above estimate for the gas exhaustion timescale is sensitive to the adopted lower mass cut-off, the metal production depends on the mass range that we directly observe. We thus suggest that, in spite of the rather constant SFR characterizing the more recent epochs, the average past SFR in NGC 1569 should have been much lower, perhaps with short intense bursts coupled with long quiescence periods.

Considering the energy input from the recent burst of SF, VB suggest that at least part of the gas currently present in NGC 1569 will be lost from the galaxy. We confirm this expectation, since our SNII rate is consistent with what found by VB. However, the burst giving birth to the stars in our observed field did not completely exhaust the gas reservoir, since the SSCs and HII regions are younger than the field. It could be that, while the burst of SF was able to initiate a general outflow of the ISM, the highest density clumps succeeded in collapsing to form stars, possibly stimulated by the SN explosions. The ultimate fate of the outflowing gas depends on several uncertain factors, including dissipation. In addition,

hydrodynamic models for bursting dwarfs suggest that, once the energy input from stellar winds and supernovae has ceased, the interstellar medium in the outer disk of the galaxy falls back to the center, where it could power a successive burst of SF (D’Ercole & Brighenti 1998).

To summarize, the recent burst could be the strongest SF episode in this galaxy over its entire lifetime, though the future evolution is rather uncertain. Among the various models proposed in the literature to explain the excess population of faint blue galaxies, we consider those by Gronwall & Koo (1995) and Babul & Rees (1992). In the model proposed by Gronwall & Koo (1995), the blue galaxy population does not evolve. The luminosity evolution of NGC 1569 is likely to be characterized by on and off phases, corresponding to the recurrent bursts, with long interburst periods. Therefore, if NGC 1569 were a typical object in the class of galaxies mostly contributing to the excess blue objects, a constant luminosity evolution of the population could be accomplished only with an appropriate number evolution of the galaxies of this kind — not a very attractive scenario. In other words, it seems unlikely that NGC 1569 represents the local counterpart of the galaxies mostly contributing to the faint blue counts in the Gronwall & Koo (1995) model.

According to Babul & Rees (1992), the excess blue objects are bursting dwarf galaxies (at redshifts close to  $\sim 1$ ) which, after the burst, become gas-free because of SN explosions. Later evolution is bound to passive fading. Thus, NGC 1569 cannot be regarded as a possible descendant of the bursting dwarfs, since it is bursting now. Still, its properties may be relevant to the Babul & Rees model. In a quantitative study of this models, Babul & Ferguson (1996) show that the current constraints on the faint galaxy population can be met if the SFR in the bursting dwarfs is large, so that the object is bright enough to contribute to the counts at the appropriate redshift. Our derived SFR for the most recent burst in NGC 1569 seems close to the rate of  $\sim 1 \text{ M}_{\odot} \text{ yr}^{-1}$  required by their model.

However, Babul & Ferguson adopt a burst duration of 10 Myr, one order of magnitude lower than what found here for NGC 1569. The success of their model depends on the combination of the duration of the burst, efficiency of star formation, and on the decay rate of the dwarf galaxy formation probability. It is likely that longer burst durations can be accommodated when varying the other parameters. The model by Babul & Ferguson also requires substantial dimming of the bursting dwarfs after the major burst. This is needed in order to avoid an excess of objects at brighter magnitudes, where the counts are dominated by the evolution of bigger galaxies. As emphasized before, we do not know if NGC 1569 will definitely lose its gas. The fact that NGC 1569 has experienced at least one previous burst and that the most recent one has had a relatively long duration, raises doubts on this hypothesis. Nevertheless, the case of NGC 1569 demonstrates that dwarf galaxies are capable to sustain bursts of SF of the magnitude required by this model.

## 7. Summary and conclusions

In this paper we have presented HST WFPC2 F555W and F439W photometry of resolved stars in the dwarf irregular galaxy NGC 1569. About 2800 stars were measured with photometric error smaller than  $\sigma_{DAO} = 0.2 \text{ mag}$  out of a total of  $\sim 7000$  objects detected. The corresponding CMD extends down to  $m_{F439}, m_{F555} \simeq 26$ , and is complete for  $m_{F555} \lesssim 23$ . It samples stars more massive than  $\sim 4 M_{\odot}$  allowing us to derive the SF history over the last 0.15 Gyr.

The interpretation of the data has been performed via theoretical simulations, based on stellar evolutionary tracks. We have considered several sets of stellar tracks from the Geneva and Padova groups, and were able to reach a satisfactory representation of the data adopting either the Geneva tracks with  $Z = 0.001$ , or the Padova tracks with  $Z = 0.004$ . Since the applicability of these tracks is related to the shape of the blue loops, which

is sensitive to details in the input parameters used in the computations, we consider it premature to dismiss alternative sets of evolutionary tracks.

The two sets provide a similar picture of the recent SF history in the field of NGC 1569. The galaxy has experienced a general burst of SF, of duration  $\Delta t \gtrsim 0.1$  Gyr, at an approximately constant rate. The burst seems to have stopped  $\sim 5 - 10$  Myr ago but SF continues in the HII regions and in the SSCs. If this general burst consists of successive episodes, these must have occurred at a similar rate, and be separated by short quiescent periods. Qualitatively, this behavior looks similar to that inferred for dwarf irregulars in the Local Group. Quantitatively, though, the level of the SFR in this recent burst in NGC 1569 is approximately 2 orders of magnitude higher.

We find that the Salpeter IMF is consistent with the observed CMD and LF, but our best simulations are characterized by slightly steeper slopes ( $2.6 - 3$ ). This seems to disagree with observations of other starbursts which generally indicate a Salpeter slope (Leitherer 1997). The present study, however, refers to the field population of stars less massive than  $\sim 30 M_{\odot}$ , whereas most starburst IMFs are derived for the nuclear burst and for dense OB clusters in the mass range above  $\sim 20 M_{\odot}$ . Since most of the SF activity in the NGC 1569 field has subsided about  $7 - 10$  Myr ago, we have no direct information on the IMF of those stars with lifetimes less than this value. The corresponding minimum masses are around  $30 M_{\odot}$ . The field-star IMF in our Galaxy and in the Magellanic Clouds is steeper (biased against massive stars) than the cluster IMF (Massey, Johnson, & DeGioia-Eastwood 1995a; Massey et al. 1995b), with the cluster value being close to Salpeter. This could be due to a richness effect, where the most massive stars are not observed simply because of their small expected numbers. NGC 1569 may be another case where the most massive stars with masses above  $\sim 30 M_{\odot}$  are preferentially formed in clusters, but we emphasize the different mass range sampled by our and that of LMC/SMC studies. Our preferred IMF

is quite similar to that derived for field stars of comparable mass in our Galaxy. This is in agreement with the notion of a virtually constant IMF, at least as far as the shape is concerned, and does not support the expectations of flatter slopes in starbursting regions (e.g., Padoan, Nordlund, & Jones 1997), or in low metallicity environments, as sometimes invoked to better reproduce the properties of elliptical galaxies (e.g., Vazdekis et al. 1996).

In the past, the SF in NGC 1569 is likely to have proceeded at a substantially lower average rate than in the recent burst. This follows from the estimate of the gas exhaustion timescale, and from the relatively low metallicity of the ISM in this galaxy. This leaves the possibilities of either an approximately constant SFR, or short strong bursts, with long interburst periods. Detailed chemical evolution models are needed to explore quantitatively the possible evolutionary paths for NGC 1569, and this will be the subject of a forthcoming paper.

The recent burst of SF is certainly not the first episode in NGC 1569 (see also VB) but may be the last. The velocity field of the ionized gas (Tomita, Ohta, & Saito 1994) and energetics arguments support this possibility (Heckman et al. 1995). If this was the case, in the future NGC 1569 may turn into a dwarf spheroidal galaxy. In addition, its luminosity evolution would be characterized by passive fading, which seems required by the Babul & Ferguson (1996) bursting dwarfs model to explain the excess faint galaxy counts. However, it is also possible that the outflowing gas falls back onto the galaxy, triggering a successive burst of SF, an option which has to be explored with the computation of detailed hydrodynamic modelling for bursting dwarf galaxies (D’Ercole & Brighenti 1998). In spite of these uncertainties, our results shows that dwarf galaxies are capable of sustaining bursts of SF at large enough rates to be relevant for the interpretation of the faint galaxy counts.

This work has been partially supported by ASI through grant 9770070AS. Support for this work was also provided by NASA grant GO-06111.01-94A from the Space Telescope



Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. L.G. and M.T. gratefully acknowledge the hospitality of STScI.

## REFERENCES

- Aparicio, A., Gallart, C., & Bertelli, G., 1997a, *AJ*, 114, 669
- . 1997b, *AJ*, 114, 680
- Arp, H., & Sandage, A. 1985, *AJ*, 90, 1163
- Babul, A., & Ferguson, H. C. 1996, *ApJ*, 458, 100
- Babul, A., & Rees, M. J. 1992, *MNRAS*, 255, 346
- Bertelli, P., Mateo, M., Chiosi, C., & Bressan, A. 1992, *ApJ*, 388, 400
- Binggeli, B. 1994, in *Dwarf Galaxies*, ed. G. Meylan & P. Prugniel (Garching: ESO), 13
- Biretta, J. 1996, *WFPC2 Instrument Handbook* (Baltimore: STScI)
- Bresolin, F., Capaccioli, M., & Piotto, G. 1993, *AJ*, 105, 1779
- Calzetti, D., Kinney, A., & Storchi-Bergmann, T. 1994, *ApJ*, 429, 582
- Charbonnel, C., Meynet, A., Maeder, A., Schaller, G., & Schaerer, D. 1993, *A&AS*, 101, 415
- D’Ercole, A., & Brighenti, F. 1998, in preparation
- Devost, D., Roy, J.-R., & Drissen, L. 1997, *ApJ*, 482, 765
- Della Ceca, R., Griffiths, R. E., Heckman, T. M., & MacKenty, J. W. 1996, *ApJ*, 469, 662
- De Marchi, G., Clampin, M., Greggio, L., Leitherer, C., Nota, A., & Tosi, M. 1997, *ApJ*, 479, L27
- De Young, D., & Heckman, T. M. 1994, *ApJ*, 431, 589
- Drissen, L., & Roy, J.-R. 1994, *PASP*, 106, 974
- Fagotto, F., Bressan, A., Bertelli, G., & Chiosi, C. 1994, *A&AS*, 105, 29
- Freedman, W. L. 1985, *ApJ*, 299, 74

———. 1988, *AJ*, 96, 1248

Gallagher III, J. S. 1996, in *From Stars to Galaxies: The Impact of Stellar Physics on Galaxy Evolution*, ed. C. Leitherer, U. Fritze-v. Alvensleben, & J. Huchra (San Francisco: ASP), 315

Gallagher III, J. S., & Hunter, D. A. 1984, *ARAA*, 22, 37

Gallagher III, J. S., Hunter, D. A., & Tutukov, A. V. 1984, *ApJ*, 284, 544

Gallart, C., Aparicio, A., Bertelli, G., & Chiosi, C. 1996a, *AJ*, 112, 1950

Gallart, C., Aparicio, A., & Vilchez, J. M. 1996b, *AJ*, 112, 1928

González-Delgado, R. M., Leitherer, C., Heckman, T. M., & Cerviño, M. 1997, *ApJ*, 483, 705

Gould, A., Bahcall, J. N., & Flynn, C. 1997, *ApJ*, 482, 913

Greggio, L. 1994, in *The Local Group: Comparative and Global Properties*, ed. A. Layden, R. C. Smith, & J. Storm (Garching: ESO), 72

Greggio, L., Marconi, G., Tosi, M., & Focardi, P. 1993, *AJ*, 105, 894 (GMTF)

Gronwall, C., & Koo, D. C. 1995, *ApJ*, 440, L1

Heckman, T. M. 1995, in *The Interplay between Massive Star Formation, the ISM, and Galaxy Evolution*, ed. D. Kunth, B. Guiderdoni, M. Heydari-Malayeri, & T. X. Thuan (Gif-sur-Yvette: Editions Frontieres), 159

Heckman, T. M., Armus, L., & Miley, G. K. 1990, *ApJS*, 74, 833

Heckman, T. M., Dahlem, M., Lehnert, M. D., Fabbiano, G., Gilmore, D., & Waller, W. H. 1995, *ApJ*, 448, 98

Ho, L. C., & Filippenko, A. V. 1996, *ApJ*, 466, L83

Hoessel, J. G. 1986, in *Luminous Stars and Associations in Galaxies*, ed. C. W. H. de Loore, A. J. Willis, & P. Laskarides (Dordrecht: Reidel), 439

- Holtzman, J., Burrows, C., Casertano, S., Hester, J., Trauger, J., Watson, A., & Worthey, G. 1995, *PASP*, 107, 1065
- Hunter, D. A., & Gallagher III, J. S. 1986, *PASP*, 98, 5
- Hunter, D. A., Gallagher, J. S., & Rautenkranz, D. 1982, *ApJS*, 49, 53
- Israel, F. P. 1988a, *A&A*, 194, 24
- . 1988b, *A&A*, 198, 109
- Johnson, H. L. 1966, *ARA&A*, 4, 193
- Kobulnicky, H. A., & Skillman, E. D. 1997, *ApJ*, 489, 636
- Kormendy, J. 1985, *ApJ*, 295, 73
- Kurucz, R. L. 1992, in *The Stellar Populations of Galaxies*, ed. B. Barbuy & A. Renzini (Dordrecht: Kluwer), 255
- Leitherer, C. 1997, in *The Stellar Initial Mass Function*, ed. G. Gilmore (San Francisco: ASP), in press
- Leitherer, C., Robert, C., & Drissen, L. 1992, *ApJ*, 401, 596
- Lejeune, T., Cuisinier, F., & Buser, E. 1997, *A&AS*, 125, 229
- Marconi, G., Matteucci, F., & Tosi, M. 1994, *MNRAS*, 270, 35
- Marconi, G., Tosi, M., Greggio, L., & Focardi, P. 1995, *AJ*, 109, 173 (MTGF)
- Marlowe, A. T. 1997, Ph.D. thesis (Baltimore: The Johns Hopkins University)
- Martin, C. L. 1997, *ApJ*, 491, 561
- Massey, P., Johnson, K. E., & DeGioia-Eastwood, K. 1995a, *ApJ*, 454, 151
- Massey, P., Lang, C. C., DeGioia-Eastwood, K., & Garmany, C. D. 1995b, *ApJ*, 438, 188
- Matteucci, F., & Tosi, M. 1985, *MNRAS*, 217, 391

- Melnick, J., Moles, M., & Terlevich, R. 1985, *A&A*, 149, L24
- Meurer, G. 1995, *Nature*, 375, 742
- Meynet, A., Maeder, A., Schaller, G., Schaerer, D., & Charbonnel, C. 1994, *A&AS*, 103, 97
- O’Connell, R. W., Gallagher, J. S., & Hunter, D. A. 1994, *ApJ*, 433, 65
- Padoan, P., Nordlund, A. P., & Jones, B. J. T. 1997, *MNRAS*, 288, 145
- Pilyugin, L. S. 1994, *A&A*, 287, 387
- Renzini, A., Greggio, L., Ritossa, C., & Ferrario, L. 1992, *ApJ*, 400, 280
- Ritossa, C. 1996, *MNRAS*, 281, 970
- Sandage, A., & Brucato, R. 1979, *AJ*, 84, 42
- Scalo, J. M. 1986, *Fund. Cosm. Phys.*, 11, 1
- Schaerer, D., Meynet, A., Maeder, A., & Schaller, G. 1993, *A&AS*, 98, 523
- Schaller, G., Schaerer, D., Meynet, A., & Maeder, A. 1992, *A&AS*, 96, 269
- Silk, J. 1987, in *High Redshift and Primeval Galaxies*, ed. J. Bergeron, D. Kunth, B. Rocca-Volmerange, & J. Tran Than Van (Gif-sur-Yvette: Editions Frontieres), 301
- Smecker-Hane, T. A., Stetson, P. B., Hesser, J. E., & Lehnert, M. D. 1994, *AJ*, 108, 507
- Suchkov, A. A., Balsara, D. S., Heckman, T. M., & Leitherer, C. 1994, *ApJ*, 430, 511
- Tammann, G. 1994, in *Dwarf Galaxies*, ed. G. Meylan & P. Prugniel, (Garching: ESO), 3
- Tenorio-Tagle, G., & Muñoz-Tuñón, C. 1997, *ApJ*, 478, 134
- Terlevich, R., Melnick, J., Masegosa, J., Moles, M., & Copetti, M. V. F. 1991, *A&AS*, 91, 285
- Thuan, T. X. 1991, in *Massive Stars in Starbursts*, ed. C. Leitherer, N. Walborn, T. Heckman, & C. Norman (Cambridge: CUP), 183

- Timmes, F. X., Woosley, S. E., & Weaver, T. A. 1995, *ApJS*, 98, 617
- Tomita, A., Ohta, K., & Saito, M. 1994, *PASJ*, 295, 73
- Tolstoy, E. 1996, *ApJ*, 462, 684
- Tolstoy, E., & Saha, A. 1996, *ApJ*, 462, 672
- Tosi, M. 1994, in *Dwarf Galaxies*, ed. G. Meylan & P. Prugniel (Garching: ESO), 465
- Tosi, M., Greggio, L., Marconi, G., & Focardi, P. 1991, *AJ*, 102, 951 (TGMF)
- Vallenari, A., & Bomans, D. J. 1996, *A&A*, 313, 713 (VB)
- van den Bergh, S. 1977, in *The Evolution of Galaxies and Stellar Populations*, ed. B. M. Tinsley & R. B. Larson (New Haven: Yale), 19
- Vazdekis, A., Casuso, E., Peletier, R. F., & Beckman, J. E. 1996, *ApJS*, 106, 307
- Walborn, N. R. 1991, in *Massive Stars in Starbursts*, ed. C. Leitherer, N. Walborn, T. M. Heckman, & C. A. Norman (Cambridge: CUP), 145
- Waller, W. 1991, *ApJ*, 370, 144
- Wirth, A., & Gallagher III, J. S. 1984, *ApJ*, 282, 85

Fig. 1.— Photometric errors versus magnitude as given by DAOPHOT in the F439W (left panel) and F555W (right panel) filters.

Fig. 2.— CMD of the stars detected in the PC field of NGC 1569: a) objects with photometric error  $<0.2$  mag, b) objects with error  $<0.1$  mag.

Fig. 3.— Luminosity function of stars with photometric error  $\sigma_{DAO} < 0.1$ . Full dots correspond to the whole set of data shown in Fig. 2 (panel b), open circles to the *reference blue plume* defined in the text.

Fig. 4.— Geneva tracks with  $Z = 0.001$  in the observational plane, having adopted  $E(B - V) = 0.56$ ,  $(m - M)_0 = 26.71$ . The faintest loop appearing in the plot belongs to the  $5 M_{\odot}$  track. The other sequences have  $M=7,9,12,15,20,25,40,60,85,120 M_{\odot}$ . Only the main-sequence evolution is plotted for the three most massive sequences. Each dot corresponds to a model in the evolutionary sequence. The slow evolutionary phases can be easily recognized from the high density of dots along each track.

Fig. 5.— Synthetic CMD computed with Geneva tracks. Panel a) shows one of the best cases obtained with the  $Z = 0.008$  set; panel b) shows one of the best results with the  $Z = 0.004$  set. The corresponding global LFs (solid line for the  $Z = 0.008$  case, dotted for the  $Z = 0.004$  case) is compared with the observational LF in panel c). See text for more details.

Fig. 6.— Geneva tracks with  $Z = 0.008$  in the observational plane, having adopted  $E(B - V) = 0.46$ ,  $(m - M)_0 = 26.71$ . The faintest loop appearing in the plot belongs to the  $5 M_{\odot}$  track. The other sequences have  $M=7,10,12,15,20,25,40,60,85,120 M_{\odot}$ . For the three most massive sequences only the Main Sequence evolution is plotted. Each dot corresponds to a model in the evolutionary sequence.

Fig. 7.— Synthetic CMD computed with Geneva tracks with  $Z = 0.001$ . Panel a) shows the best result obtained with constant SF, panel b) shows the best result obtained with an exponentially decreasing SF. Both simulations assume no SF activity during the last 7 Myr. The corresponding global LF (solid line for the panel a model, dotted for the panel b model) is compared with the observational LF in panel c.

Fig. 8.— Synthetic CMD computed with Geneva tracks with  $Z = 0.001$ . Panels a) and b) show the best results obtained with two distinct episodes of SF, the older one ending 0.05 Gyr ago. Compared to the case shown in panel b), in the simulation shown in panel a) we have adopted a larger SFR in the most recent episode. This trend is counterbalanced by the different IMF exponent. In panel c) we show the luminosity functions for the models in panel a) (solid line) and panel b) (dotted line), and the observed LF (dots).

Fig. 9.— Synthetic CMD computed with the Padova tracks with  $Z=0.004$ . Panels a) and b) show the best results obtained with two episodes of SF, the older one ending 0.035 Gyr ago. In panel c) we show the luminosity functions for the models in panel a) (solid line) and panel b) (dotted line). In both cases the SFR in the most recent episode is slightly lower than in the previous one, and one can see the effect of varying the IMF exponent: the steeper slope (panel b)) tends to underpopulate the brightest bins of the LF. No SF is assumed to have occurred in the last 10 Myr.

Fig. 10.— Synthetic CMD computed with the Geneva tracks with  $Z=0.001$ , a constant SF and  $\alpha=3$ , but adopting different epochs for the end of the burst: currently ongoing SF (panel a), stopped at  $5 M_{\odot} \text{ yr}^{-1}$  ago (panel b). The corresponding global LF (solid line for the panel a) model, dotted for the panel b) model) is compared with the observational LF in panel c).

Fig. 11.— Synthetic CMD computed with the Padova tracks with  $Z=0.004$ ,  $(m - M)_0 =$



26.71 and  $E(B - V)=0.46$ . The synthetic CMD in panel a) has been obtained assuming two episodes of SF, separated by a quiescence period of 10 Myr (from 40 to 30 Myr ago), and no SF in the last 10 Myr. In the simulation in panel b) a constant, currently ongoing SF has been adopted. The corresponding LFs (solid line for the CMD in the top panel) are shown in panel c).

Fig. 12.— Synthetic CMD computed with the Padova tracks with  $Z=0.004$ , and adopting  $(m - M)_0 = 28.01$  and  $E(B - V)=0.46$ . The simulation in panel a) has been computed assuming two episodes of SF, the most recent one stopped 5 Myr ago. In the simulation in panel b) a currently ongoing, constant SFR has been adopted. As usual, panel c) shows the corresponding LFs (solid line for the CMD in the top panel).

Table 1: Photometric completeness as a function of the magnitude in the F439W band

| $m_{439}$     | completeness  |
|---------------|---------------|
| $< 23.0$      | $> 99\%$      |
| $23.0 - 24.0$ | $\simeq 80\%$ |
| $24.0 - 25.0$ | $\simeq 50\%$ |
| $25.0 - 26.0$ | $\simeq 30\%$ |

Table 2: Stellar evolutionary tracks adopted for the synthetic CMDs

| Model  | $Z$   | $\dot{M}$ | Reference                |
|--------|-------|-----------|--------------------------|
| Geneva | 0.008 | standard  | Schaerer et al. (1993)   |
| Geneva | 0.008 | high      | Meynet et al. (1993)     |
| Geneva | 0.004 | standard  | Charbonnel et al. (1993) |
| Geneva | 0.004 | high      | Meynet et al. (1993)     |
| Geneva | 0.001 | standard  | Schaller et al. (1992)   |
| Padova | 0.004 | standard  | Fagotto et al. (1994)    |

























